
The chromatin regulator Brg1 suppresses formation of intraductal papillary mucinous neoplasm and pancreatic ductal adenocarcinoma.

Journal: Nat Cell Biol

Publication Year: 2014

Authors: Guido von Figura, Akihisa Fukuda, Nilotpal Roy, Muluye E Liku, John P Morris Iv, Grace E Kim, Holger A Russ, Matthew A Firpo, Sean J Mulvihill, David W Dawson, Jorge Ferrer, William F Mueller, Anke Busch, Klemens J Hertel, Matthias Hebrok

PubMed link: 24561622

Funding Grants: Training Program in Stem Cell Research at UCSF

Public Summary:

Scientific Abstract:

Pancreatic ductal adenocarcinoma (PDA) develops through distinct precursor lesions, including pancreatic intraepithelial neoplasia (PanIN) and intraductal papillary mucinous neoplasia (IPMN). However, genetic features resulting in IPMN-associated PDA (IPMN-PDA) versus PanIN-associated PDA (PanIN-PDA) are largely unknown. Here we find that loss of Brg1, a core subunit of SWI/SNF chromatin remodelling complexes, cooperates with oncogenic Kras to form cystic neoplastic lesions that resemble human IPMN and progress to PDA. Although Brg1-null IPMN-PDA develops rapidly, it possesses a distinct transcriptional profile compared with PanIN-PDA driven by mutant Kras and hemizygous p53 deletion. IPMN-PDA also is less lethal, mirroring prognostic trends in PDA patients. In addition, Brg1 deletion inhibits Kras-dependent PanIN development from adult acinar cells, but promotes Kras-driven preneoplastic transformation in adult duct cells. Therefore, this study implicates Brg1 as a determinant of context-dependent Kras-driven pancreatic tumorigenesis and suggests that chromatin remodelling may underlie the development of distinct PDA subsets.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/chromatin-regulator-brg1-suppresses-formation-intraductal-papillary-mucinous>